



# The Laplacian Energy of an Intuitionistic Fuzzy Rough Graph and its Utilisation in Decision-Making

Shaik Noorjahan<sup>1</sup> Shaik Sharief Basha<sup>\*2</sup>

<sup>1</sup>Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology, Vellore, Tamil Nadu, India.

<sup>2</sup>Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology, Vellore, Tamil Nadu, India.

\*Corresponding author; email address: [shariefbasha.s@vit.ac.in](mailto:shariefbasha.s@vit.ac.in)

## Abstract

An intuitionistic fuzzy rough model is a hybrid model that combines intuitionistic fuzzy sets and rough sets, addressing soft computing and ambiguity. It uses lower and upper approximation spaces in various fields, including science, technology, database systems, computer networks, and expert system architecture. The matrix of adjacency of an intuitionistic fuzzy rough graph is described in the article. The matrix of adjacency also provides the upper and lower bounds for Laplacian energy. These are used to define the Laplacian energies of intuitionistic fuzzy rough graphs and the weight function of Laplacian energy within these graphs. The intuitionistic fuzzy rough preference relation method is used to process the intuitionistic fuzzy rough weighted average. This technique is applied in data visualizations, which make data understandable by displaying it in a graphical or pictorial style. It supports decision-making and provides factual justification. This approach benefits any field that needs creative methods for presenting large volumes of complex data. Modern computer graphics have significantly influenced visualization.

**Keywords:** Laplacian energy (LE), intuitionistic fuzzy rough graph (IFRG), data visualization, and decision-making

## 1. Introduction

*Graph theory*, developed by Euler in 1736, is a versatile tool used in various disciplines, including geometry, transportation, and engineering. Fuzzy sets and fuzzy logic were introduced by Zadeh in 1965 [32], convey ambiguity and uncertainty, and are studied in various disciplines like medicine, management, and artificial intelligence. A new degree, known as the degree of non-membership, was introduced to the fuzzy sets idea by Atanassov [4] in 1986 to resolve the ambiguity and uncertainty around the membership degree. Rosenfeld [25] pioneered fuzzy graphs, a concept first introduced by Kaufmann [17] in 1973. He proposed fuzzy relations and developed the basic framework of fuzzy graphs. Atanassov [3] is credited with the development of intuitionistic fuzzy graphs and fuzzy relations.

Jamil et al. [16] uses Sombor indices to study intuitionistic fuzzy graphs, which are generalizations of fuzzy graphs, to assess vaccination centres during the pandemic, highlighting their practical implications and efficiency. Pawlak's [21] rough set theory is considered innovative in soft processing tools as it uniquely addresses ambiguity. Rough set theory uses lower and upper approximation ideas to extract data from systems of data and represent it as decision rules, addressing ambiguity in the universe. In Chakrabarty et al. [7], the fuzziness of imperfect sets was explored. The concept of a fuzzy rough set [11] has been developed to enhance decision-making efficiency. This paper [14] explores rough graphs for uncertainty problems, class connections, subgraphs, weighted rough graphs, and an algorithm for exploring class optimal trees in weighted rough graphs. Akram and Arshad [1] introduced fuzzy rough graphs with application.

Intuitionistic fuzzy rough sets were described by Cornelis et al. [9]. The intuitionistic fuzzy rough sets model based on operators was investigated by Xu et al. [31]. Types of intuitionistic fuzzy rough approximation operations are studied by Zhou and Wu [35], while Haq et al. [13] tackle semantic issues with incomplete information by classifying types, proposing a fuzzy decision table, and developing rule extraction methods. Zhang and Zhu [34] developed three asymmetric models for resolving uncertain problems in an intuitionistic fuzzy environment, combining traditional PROMETHEE and TOPSIS procedures, demonstrating their effectiveness. Tiwari and Lohani [28] introduce a novel Interval-valued intuitionistic fuzzy rough set (IVIFRS) system for conflict analysis in decision-making, utilizing rough set and interval-valued intuitionistic fuzzy set theories to measure disputes and identify conflicting attributes. Hussain et al. [15] introduce a TOPSIS method utilizing Dombi operations for aggregating averaging and geometric operators, specifically for MCGDM, specifically designed for diagnosing severe COVID-19 patients. With an emphasis on fuzzy preference relations for effective alternative assessment, comparison, selection, and ordering, Borzecka [6] investigates fuzzy multi-criteria decision-making utilizing Zadeh's linguistic method. In order to compare criterion significance, prove compositional qualities and novel relational features, provide consistent choices, and rank options, Peneva and Popchev [22] investigates the weighted aggregation of fuzzy preference relations for decision-making issues.

Akram et al. [19, 33] introduced several innovative ideas in intuitionistic fuzzy rough graphs for decision-making. This [2] monograph explores hybrid models, specifically fuzzy digraphs, for complex problem-solving, highlighting their applications in decision-making and overcoming the uncertainty issues of traditional methods. Mahmood et al. [18] and Mazarbhuiya and Shenify [20] present innovative techniques for precise disease diagnosis and anomaly detection, combining rough set theory and intuitionistic fuzzy set theory for high accuracy rates. Tiwari [29] introduces a novel intuitionistic fuzzy (IF)-assisted mutual information concept that effectively handles noise, uncertainty, and vagueness in real-valued datasets, improving phospholipids positive molecule prediction. Salamat et al. [26] describe the utilization of intuitionistic fuzzy sets and rough sets in informal computing for sparse data, aiming for accuracy, accessibility, and cost-effectiveness.

Gutman introduced the concept of graph energy in chemistry, discovering lower and upper bounds on the total electron energy of certain molecules. Chemistry's relationship between molecular graph energy and overall electron energy is crucial, as a graph with  $n$  nodes has energy  $2(n - 1)$ , while a graph with separated vertices has energy zero. Gutman and Zhou [12] defined the Laplacian energy (LE) of a graph as the sum of the absolute values of discrepancies between the average vertex degree and its

Laplacian eigenvalues. Rahimi Sharbaf and Fayazi [24] defined a fuzzy graph's Laplacian energy. Basha and Kartheek [5] generalised the Laplacian energy of an intuitionistic fuzzy graph from the Laplacian energy of a fuzzy graph. Poonia and Bajaj [23] introduce a decision-making methodology for ranking alternatives in fuzzy graphs or directed graphs for hydropower plant site selection, analyzing its originality, comparative remarks, advantageous features, and limitations. Sarkar introduced advanced software to the pharmaceutical industry for tasks such as record-keeping, quality control, and clinical trial data management [27]. Dave et al. provide a concise discussion on data visualization, highlighting various tools and techniques for effectively communicating insights [10].

### 1.1. Challenges and Motivation

Based on the overhead review, it can be concluded that an intuitionistic fuzzy rough graph (IFRG) aims to provide a reliable and adaptable mathematical framework for analyzing systems under imprecision and uncertainty in practical applications. IFRGs are also often used to reflect uncertainty in decision problems. Therefore, combinations of this tool with multi-criteria decision-making (MCDM) methods are continuously being developed. Due to the difficulty of choosing the proper MCDM techniques, further work on their development should be carried out. Moreover, a frequently addressed issue in IFRGs is the aggregation of expert knowledge. Numerous aggregation operators need to be continuously investigated.

Given the above constraints associated with IFRGs, the motivation of this study is as follows:

1. There is a lack of comparative analysis between the results obtained from IFRGs in some works;
2. Specific terms concentrate on a specific purpose that combines domain expertise;
3. Works considering IFRGs in real-life problems are missing;
4. Few studies consider using MCDM techniques in conjunction with IFRGs.

### 1.2. Contributions and Novelties

This paper focuses on using the Laplacian energy of an intuitionistic fuzzy rough graph approach in a data visualization problem. Laplacian energy is used to determine decision weights based on decision-maker's preferences. Three decision-makers were involved in the whole process and presented their strategies using intuitionistic fuzzy rough graph. Then, evaluate the aggregated preferences of the decision-makers. These preferences were also aggregated into four functions.

### 1.3. Framework of the study

In the research paper, organized as follows: Section 2 contains fundamental concepts related to intuitionistic fuzzy rough graphs. In Section 3, we describe an intuitionistic fuzzy rough graph's Laplacian energy with example. In Section 4, the lower and upper bounds for an intuitionistic fuzzy rough graph's Laplacian energy are also determined by its properties. The Laplacian energy of an intuitionistic fuzzy rough digraph is described in Section 5. Data visualization serves as a real-world example, and we use a comparison study to demonstrate these concepts in Section 6. Finally, Section 7 presents conclusions and future research directions.

## 2. Preliminaries

**Definition 1.** [1] Considering that  $F$  is an ordered tuple  $(P_i, P_iR, Q_i, Q_iS)$ , a non-empty set with a fuzzy rough graph of  $G$ .

1.  $P_i$  denotes a fuzzy relation to  $F$ ,
2.  $Q_i$  denotes a fuzzy relation to  $D \subseteq F \times F$ ,
3.  $P_iR = (\underline{P_iR}, \overline{P_iR})$  denotes a fuzzy rough set to  $F$ ,
4.  $Q_iS = (\underline{Q_iS}, \overline{Q_iS})$  denotes a fuzzy rough relation to  $F$ .
5. Therefore,  $G = ((\underline{P_iR}, \overline{P_iR}), (\underline{Q_iS}, \overline{Q_iS}))$  are lower and upper approximations on fuzzy rough graphs of  $G, \forall c, d \in F$ .

$$(\underline{Q_iS})(cd) \leq \min \{(\underline{P_iR})(c), (\underline{P_iR})(d)\}, (\overline{Q_iS})(cd) \leq \min \{(\overline{P_iR})(c), (\overline{P_iR})(d)\}.$$

**Definition 2.** [2] A four-ordered tuple  $(X_i, X_iU, Y_i, Y_iV)$  is an intuitionistic fuzzy rough graph  $G$  on a nonempty set  $F$ .

1.  $X_i$  represents an intuitionistic fuzzy relation to  $F$ ,
2.  $Y_i$  represents an intuitionistic fuzzy relation to  $D \subseteq F \times F$ ,
3.  $X_iU = (\underline{X_iU}, \overline{X_iU})$  represents an intuitionistic fuzzy rough set to  $F$ ,
4.  $Y_iV = (\underline{Y_iV}, \overline{Y_iV})$  represents an intuitionistic fuzzy rough set to  $F$ .
5. Thus  $G = (\underline{G}, \overline{G}) = (X_iU, Y_iV)$  is an intuitionistic fuzzy rough graph. Here  $\underline{G} = (\underline{X_iU}, \overline{X_iU})$  and  $\overline{G} = (\underline{Y_iV}, \overline{Y_iV})$  are lower and upper approximations of the intuitionistic fuzzy rough graph of  $G, \forall l, m \in F$ .

$$(\underline{Y_iV})^+(lm) \leq \min \{(\underline{X_iU})^+(l), (\underline{X_iU})^+(m)\}, (\underline{Y_iV})^-(lm) \leq \max \{(\underline{X_iU})^-(l), (\underline{X_iU})^-(m)\},$$

$$(\overline{Y_iV})^+(lm) \leq \min \{(\overline{X_iU})^+(l), (\overline{X_iU})^+(m)\}, (\overline{Y_iV})^-(lm) \leq \max \{(\overline{X_iU})^-(l), (\overline{X_iU})^-(m)\}.$$

## 3. Laplacian energy of an intuitionistic fuzzy rough graph

**Definition 3.** The matrix of adjacency  $A(\xi) = (A(\mu_N(p_i p_j)), A(\nu_N(p_i p_j)))$  of an IFRG  $\xi = (\underline{\xi}, \overline{\xi}) = (M, N)$  is defined as a square matrix  $A(\xi) = [a_{ij}]$ , where  $a_{ij} = (\mu_N(p_i p_j), \nu_N(p_i p_j))$ .

Here,  $\mu_N(p_i p_j), \nu_N(p_i p_j)$  represent the degree of the relationship and degree of non-relationship between  $p_i$  and  $p_j$ , respectively.

**Definition 4.** Consider  $p_i$  as a vertex of the intuitionistic fuzzy rough graph  $\xi = (\underline{\xi}, \overline{\xi}) = (M, N)$ . The degree of  $u$  is denoted as  $d_\xi(p_i) = \sum_{p_i p_j \in E(\xi)} \mu(p_i p_j)$ .

**Definition 5.** Let  $\xi = (\underline{\xi}, \overline{\xi}) = (M, N)$  be an IFRG on  $n$  vertices. The degree matrix,

$D(\xi) = (D(\mu_N(p_i p_j)), D(\nu_N(p_i p_j))) = [d_{ij}]$ , of  $\xi$  is a  $n \times n$  diagonal matrix defined as:

$$d_{ij} = \begin{cases} d_\xi(p_i) & \text{if } i = j; \\ 0 & \text{otherwise.} \end{cases}$$

**Definition 6.** The Laplacian matrix of an IFRG  $\xi = (\underline{\xi}, \overline{\xi}) = (M, N)$  is defined as

$L(\xi) = (L(\mu_N(p_i p_j)), L(\nu_N(p_i p_j))) = D(\xi) - A(\xi)$ , where  $A(\xi)$  is an adjacency matrix and  $D(\xi)$  is a

degree matrix of an IFRG  $\xi$ .

**Definition 7.** The Laplacian matrix of spectrum of an IFRG  $L(\xi)$  is denoted by  $(R_L, S_L)$ , here  $R_L = L(\mu_N(p_i p_j))$  and  $S_L = L(\nu_N(p_i p_j))$  are the sets of Laplacian eigenvalues.

**Definition 8.** The Laplacian energy of an IFRG  $\xi = (\underline{\xi}, \bar{\xi}) = (M, N)$  is defined as:

$$LE(\underline{\xi}) = (LE(\underline{\mu}_N(p_i p_j)), LE(\underline{\nu}_N(p_i p_j))) = \left( \sum_{i=1}^n |\underline{\rho}_i|, \sum_{i=1}^n |\underline{\sigma}_i| \right),$$

$$LE(\bar{\xi}) = (LE(\bar{\mu}_N(p_i p_j)), LE(\bar{\nu}_N(p_i p_j))) = \left( \sum_{i=1}^n |\bar{\rho}_i|, \sum_{i=1}^n |\bar{\sigma}_i| \right).$$

Here:

$$\underline{\rho}_i = \underline{\phi}_i - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n}, \quad \underline{\sigma}_i = \underline{\psi}_i - \frac{2 \sum_{1 \leq i < j \leq n} \nu_N(p_i p_j)}{n},$$

$$\bar{\rho}_i = \bar{\phi}_i - \frac{2 \sum_{1 \leq i < j \leq n} \bar{\mu}_N(p_i p_j)}{n}, \quad \bar{\sigma}_i = \bar{\psi}_i - \frac{2 \sum_{1 \leq i < j \leq n} \bar{\nu}_N(p_i p_j)}{n}.$$

**Definition 9.** If the weight  $w_r = (\underline{w}_r, \bar{w}_r)$  of a wide range of Laplacian energy of an intuitionistic fuzzy rough graph is calculated using the formula below.

For lower,

$$\underline{w}_r = \left( (\underline{w}_\mu)_r, (\underline{w}_\nu)_r \right) = \left( \frac{LE((\underline{D}_\mu)_r)}{\sum_{i=1}^s LE((\underline{D}_\mu)_i)}, \frac{LE((\underline{D}_\nu)_r)}{\sum_{i=1}^s LE((\underline{D}_\nu)_i)} \right),$$

For upper,


$$\bar{w}_r = \left( (\bar{w}_\mu)_r, (\bar{w}_\nu)_r \right) = \left( \frac{LE((\bar{D}_\mu)_r)}{\sum_{i=1}^s LE((\bar{D}_\mu)_i)}, \frac{LE((\bar{D}_\nu)_r)}{\sum_{i=1}^s LE((\bar{D}_\nu)_i)} \right),$$

$r = 1, 2, \dots, s$ .

**Example 1.** Let a graph  $G = (V, E)$ . where  $V = \{p_1, p_2, p_3, p_4, p_5\}$  and  $E = \{p_1 p_2, p_2 p_3, p_3 p_4, p_4 p_5, p_5 p_1, p_1 p_3, p_1 p_4, p_2 p_4, p_2 p_5\}$ . Let  $\xi = (\underline{\xi}, \bar{\xi}) = (M, N)$  be an IFRG on  $V$  as show in Figure (1).

The adjacency matrix, degree matrix and Laplacian matrix of a lower IFRG shown in Figure (1a) are as follows:

$$A(\underline{\xi}) = \begin{bmatrix} (0.0, 0.0) & (0.4, 0.2) & (0.3, 0.2) & (0.4, 0.2) & (0.3, 0.2) \\ (0.4, 0.2) & (0.0, 0.0) & (0.3, 0.2) & (0.4, 0.2) & (0.3, 0.2) \\ (0.3, 0.2) & (0.3, 0.2) & (0.0, 0.0) & (0.4, 0.1) & (0.3, 0.2) \\ (0.4, 0.2) & (0.4, 0.2) & (0.4, 0.1) & (0.0, 0.0) & (0.3, 0.2) \\ (0.3, 0.2) & (0.3, 0.2) & (0.3, 0.2) & (0.3, 0.2) & (0.0, 0.0) \end{bmatrix}$$



**Figure 1.** Intuitionistic fuzzy rough graph

$$D(\underline{\xi}) = \begin{bmatrix} (1.4, 0.8) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (1.4, 0.8) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (1.3, 0.7) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (1.5, 0.7) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (1.2, 0.8) \end{bmatrix}$$

$$L(\underline{\xi}) = \begin{bmatrix} (1.4, 0.8) & (-0.4, -0.2) & (-0.3, -0.2) & (-0.4, -0.2) & (-0.3, -0.2) \\ (-0.4, -0.2) & (1.4, 0.8) & (-0.3, -0.2) & (-0.4, -0.2) & (-0.3, -0.2) \\ (-0.3, -0.2) & (-0.3, -0.2) & (1.3, 0.7) & (-0.4, -0.1) & (-0.3, -0.2) \\ (-0.4, -0.2) & (-0.4, -0.2) & (-0.4, -0.1) & (1.5, 0.7) & (-0.3, -0.2) \\ (-0.3, -0.2) & (-0.3, -0.2) & (-0.3, -0.2) & (-0.3, -0.2) & (1.2, 0.8) \end{bmatrix}$$

The Laplacian spectrum and the Laplacian energy of a lower IFRG  $\underline{\xi}$

$$Laplacian\ Spec(\underline{\mu}_N(p_i p_j)) = (0.0000, 1.5000, 1.6000, 1.8000, 1.9000),$$

$$Laplacian\ Spec(\underline{\nu}_N(p_i p_j)) = (0.0000, 0.8000, 1.0000, 1.0000, 1.0000).$$

Therefore,

$$Laplacian\ Spec(\underline{\xi}) = ((0, 0), (1.5000, 0.8000), (1.6000, 1.0000), (1.8000, 1.0000), (1.9000, 1.0000)).$$

Now,

$$LE(\underline{\xi}) = (LE(\underline{\mu}_N(p_i p_j)), LE(\underline{\nu}_N(p_i p_j))) = (2.7200, 1.5200).$$

The adjacency matrix, degree matrix and Laplacian matrix of an upper IFRG shown in Figure (1b) are as follows:

$$\begin{aligned}
A(\bar{\xi}) &= \begin{bmatrix} (0.0, 0.0) & (0.4, 0.0) & (0.5, 0.0) & (0.6, 0.0) & (0.6, 0.0) \\ (0.4, 0.0) & (0.0, 0.0) & (0.4, 0.6) & (0.5, 0.2) & (0.5, 0.1) \\ (0.5, 0.0) & (0.4, 0.6) & (0.0, 0.0) & (0.6, 0.0) & (0.6, 0.0) \\ (0.6, 0.0) & (0.5, 0.2) & (0.6, 0.0) & (0.0, 0.0) & (0.5, 0.1) \\ (0.6, 0.0) & (0.5, 0.1) & (0.6, 0.0) & (0.5, 0.1) & (0.0, 0.0) \end{bmatrix} \\
D(\bar{\xi}) &= \begin{bmatrix} (2.1, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (1.8, 0.9) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (2.1, 0.6) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (2.2, 0.3) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (2.2, 0.2) \end{bmatrix} \\
L(\bar{\xi}) &= \begin{bmatrix} (2.1, 0.0) & (-0.4, -0.0) & (-0.5, -0.0) & (-0.6, -0.0) & (-0.6, -0.0) \\ (-0.4, -0.0) & (1.8, 0.9) & (-0.4, -0.6) & (-0.5, -0.2) & (-0.5, -0.1) \\ (-0.5, -0.0) & (-0.4, -0.6) & (2.1, 0.6) & (-0.6, -0.0) & (-0.6, -0.0) \\ (-0.6, -0.0) & (-0.5, -0.2) & (-0.6, -0.0) & (2.2, 0.3) & (-0.5, -0.1) \\ (-0.6, -0.0) & (-0.5, -0.1) & (-0.6, -0.0) & (-0.5, -0.1) & (2.2, 0.2) \end{bmatrix}
\end{aligned}$$

The Laplacian spectrum and the Laplacian energy of an upper IFRG  $\bar{\xi}$

$$\text{Laplacian Spec}(\bar{\mu}_N(p_i p_j)) = (0.0000, 2.2298, 2.6000, 2.7000, 2.8702),$$

$$\text{Laplacian Spec}(\bar{\nu}_N(p_i p_j)) = (0.0000, 0.0000, 0.2284, 0.3766, 1.3950).$$

Therefore,

$$\text{Laplacian Spec}(\bar{\xi}) = ((0, 0), (2.2298, 0.0000), (2.6000, 0.2284), (2.7000, 0.3766), (2.8702, 1.3950)).$$

Now,

$$L\bar{E}(\bar{\xi}) = (L\bar{E}(\bar{\mu}_N(p_i p_j)), L\bar{E}(\bar{\nu}_N(p_i p_j))) = (4.1600, 1.9900).$$

The Laplacian energy of an intuitionistic fuzzy rough graph is

$$LE(\xi) = (LE(\underline{\xi}), L\bar{E}(\bar{\xi})) = ((2.7200, 1.5200), (4.1600, 1.9900)).$$

## 4. Results

**Theorem 1.** Let  $\xi = (\underline{\xi}, \bar{\xi}) = (M, N)$  be an IFRG, and let  $L(\xi)$  be the Laplacian matrix  $\xi$ . If  $\eta_1 \geq \eta_2 \geq \eta_3 \geq \dots \geq \eta_n$  and  $\theta_1 \geq \theta_2 \geq \theta_3 \geq \dots \geq \theta_n$  are the eigenvalues of  $L(\mu_N(p_i p_j))$  and  $L(\nu_N(p_i p_j))$ , respectively, then:

$$(i) \quad \sum_{i=1, \eta_i \in R_L}^n \eta_i = 2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j); \quad \sum_{i=1, \theta_i \in S_L}^n \theta_i = 2 \sum_{1 \leq i < j \leq n} \nu_N(p_i p_j).$$

$$(ii) \quad \sum_{i=1, \eta_i \in R_L}^n \eta_i^2 = 2 \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \sum_{i=1}^n d_{\mu_N(p_i p_j)}^2(p_i);$$

$$\sum_{i=1, \theta_i \in S_L}^n \theta_i^2 = 2 \sum_{1 \leq i < j \leq n} (\nu_N(p_i p_j))^2 + \sum_{i=1}^n d_{\nu_N(p_i p_j)}^2(p_i).$$

**Proof.** (i) Since  $L(\xi)$  contains non-negative Laplacian eigenvalues and is a symmetric matrix, such that:

$$\sum_{i=1, \eta_i \in R_L}^n \eta_i = \text{tr}(L(\xi)) = \sum_{i=1}^n d_{\mu_N(p_i p_j)}(p_i) = 2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j).$$

So,

$$\sum_{i=1, \eta_i \in R_L}^n \eta_i = 2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j).$$

Similarly, we have to prove

$$\sum_{i=1, \theta_i \in S_L}^n \theta_i = 2 \sum_{1 \leq i < j \leq n} \nu_N(p_i p_j).$$

(ii) According to the definition of a Laplacian matrix, we have:

$$L(\mu_N(p_i p_j)) = \begin{bmatrix} d_{\mu_N(p_i p_j)}(p_1) & -\mu_N(p_1 p_2) & \cdots & -\mu_N(p_1 p_n) \\ -\mu_N(p_2 p_1) & d_{\mu_N(p_i p_j)}(p_2) & \cdots & -\mu_N(p_2 p_n) \\ \vdots & \vdots & \ddots & \vdots \\ -\mu_N(p_n p_1) & -\mu_N(p_n p_2) & \cdots & d_{\mu_N(p_i p_j)}(p_n) \end{bmatrix}$$

According to the trace properties of a matrix, we have:

$$\text{tr}((L(\mu_N(p_i p_j)))^2) = \sum_{i=1, \eta_i \in R_L}^n \eta_i^2.$$

Where:

$$\begin{aligned} \text{tr}((L(\mu_N(p_i p_j)))^2) &= (d_{\mu_N(p_i p_j)}^2(p_1) + \mu_N^2(p_1 p_2) + \cdots + \mu_N^2(p_1 p_n)) + (\mu_N^2(p_2 p_1) + d_{\mu_N(p_i p_j)}^2(p_2) + \\ &\cdots + \mu_N^2(p_2 p_n)) + \cdots + (\mu_N^2(p_n p_1) + \mu_N^2(p_n p_2) + \cdots + d_{\mu_N(p_i p_j)}^2(p_n)). = 2 \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \\ &\sum_{i=1}^n d_{\mu_N(p_i p_j)}^2(p_i). \end{aligned}$$

Therefore,

$$\sum_{i=1, \eta_i \in R_L}^n \eta_i^2 = 2 \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \sum_{i=1}^n d_{\mu_N(p_i p_j)}^2(p_i),$$

Likewise, we have to prove that

$$\sum_{i=1, \theta_i \in S_L}^n \theta_i^2 = 2 \sum_{1 \leq i < j \leq n} (\nu_N(p_i p_j))^2 + \sum_{i=1}^n d_{\nu_N(p_i p_j)}^2(p_i).$$

□

**Theorem 2.** Let  $\xi = (\underline{\xi}, \bar{\xi}) = (M, N)$  be an IFRG, and let  $L(\xi)$  be the Laplacian matrix of  $\xi$ . If  $\eta_1 \geq$



$\eta_2 \geq \eta_3 \geq \dots \geq \eta_n$  and  $\theta_1 \geq \theta_2 \geq \theta_3 \geq \dots \geq \theta_n$  are the eigenvalues of  $L(\mu_N(p_i p_j))$  and  $L(\nu_N(p_i p_j))$  respectively, and  $\rho_i = \eta_i - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n}$ ,  $\sigma_i = \theta_i - \frac{2 \sum_{1 \leq i < j \leq n} \nu_N(p_i p_j)}{n}$ , then:

$$(i) \sum_{i=1}^n \rho_i = 0, \sum_{i=1}^n \sigma_i = 0;$$

$$(ii) \sum_{i=1}^n \rho_i^2 = 2P_\mu, \sum_{i=1}^n \sigma_i^2 = 2P_\nu.$$

Where:

$$P_\mu = \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \frac{1}{2} \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2,$$

$$P_\nu = \sum_{1 \leq i < j \leq n} (\nu_N(p_i p_j))^2 + \frac{1}{2} \sum_{i=1}^n \left( d_{\nu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \nu_N(p_i p_j)}{n} \right)^2.$$

**Example 1.** Consider IFRG,  $\xi = (\underline{\xi}, \bar{\xi}) = (M, N)$  and  $V = \{p_1, p_2, p_3, p_4, p_5\}$  as shown in Figure (1). Then:

From theorem (2), we have lower approximations as follows:

$$(i) \sum_{i=1}^5 \rho_i = 0, \sum_{i=1}^5 \sigma_i = 0.$$

$$(ii) \sum_{i=1}^5 \rho_i^2 = 2.4120 = 2(1.2060) = 2P_\mu, \sum_{i=1}^5 \sigma_i^2 = 0.7520 = 2(0.3760) = 2P_\nu.$$

From theorem (2), we have upper approximations as follows:

$$(i) \sum_{i=1}^5 \bar{\rho}_i = 0, \sum_{i=1}^5 \bar{\sigma}_i = 0.$$

$$(ii) \sum_{i=1}^5 \bar{\rho}_i^2 = 5.6280 = 2(2.8140) = 2\bar{P}_\mu, \sum_{i=1}^5 \bar{\sigma}_i^2 = 1.3400 = 2(0.6700) = 2\bar{P}_\nu.$$

**Theorem 3.** Let  $\xi = (\underline{\xi}, \bar{\xi}) = (M, N)$  be an IFRG on  $n$  vertices, let  $L(\xi) = (L(\mu_N(p_i p_j)), L(\nu_N(p_i p_j)))$  be the Laplacian matrix of  $\xi$ . Then,

$$(i) L\underline{E}(\mu_N(p_i p_j)), L\bar{E}(\mu_N(p_i p_j)) \leq \sqrt{2n \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + n \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2},$$

$$(ii) L\bar{E}(\nu_N(p_i p_j)), L\bar{E}(\nu_N(p_i p_j)) \leq \sqrt{2n \sum_{1 \leq i < j \leq n} (\nu_N(p_i p_j))^2 + n \sum_{i=1}^n \left( d_{\nu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \nu_N(p_i p_j)}{n} \right)^2}.$$

**Proof.** Apply Cauchy-Schwarz inequality to  $n$  numbers  $(1, 1, \dots, 1)$  and  $(|\rho_1|, |\rho_2|, \dots, |\rho_n|)$ ,

$$\sum_{i=1}^n |\rho_i| \leq \sqrt{n} \sqrt{\sum_{i=1}^n |\rho_i|^2}$$

$$L\bar{E}(\mu_N(p_i p_j)) \leq \sqrt{n} \sqrt{2P_\mu} = \sqrt{2nP_\mu}.$$

Since

$$P_\mu = \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \frac{1}{2} \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2,$$

Therefore,

$$L\bar{E}(\mu_N(p_i p_j)) \leq \sqrt{2n \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + n \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2}.$$

Similarly, we have to prove  $L\bar{E}(\mu_N(p_i p_j)), L\bar{E}(\nu_N(p_i p_j)), L\bar{E}(\nu_N(p_i p_j))$ .  $\square$

**Example 2.** Consider IFRG,  $\xi = (\xi, \bar{\xi}) = (M, N)$  and  $V = \{p_1, p_2, p_3, p_4, p_5\}$  as shown in Figure (1). Then:

From theorem (3), we have a lower approximation as follows:

$$L\bar{E}(\underline{\mu}_N(p_i p_j)) \leq \sqrt{2(5)(1.1800) + 5(0.0520)} \text{ implies } 2.7200 \leq 3.4728.$$

$$L\bar{E}(\underline{\nu}_N(p_i p_j)) \leq \sqrt{2(5)(0.3700) + 5(0.0120)} \text{ implies } 1.5200 \leq 1.9391.$$

From theorem (3), we have an upper approximation as follows:

$$L\bar{E}(\bar{\mu}_N(p_i p_j)) \leq \sqrt{2(5)(2.7600) + 5(0.1080)} \text{ implies } 4.1600 \leq 5.3047.$$

$$L\bar{E}(\bar{\nu}_N(p_i p_j)) \leq \sqrt{2(5)(0.4200) + 5(0.5000)} \text{ implies } 1.9900 \leq 2.5884.$$

**Theorem 4.** Let  $\xi = (\xi, \bar{\xi}) = (M, N)$  be an IFRG on  $n$  vertices, let  $L(\xi) = (L(\mu_N(p_i p_j)), L(\nu_N(p_i p_j)))$  be the Laplacian matrix of  $\xi$ . Then:

$$(i) L\bar{E}(\mu_N(p_i p_j)), L\bar{E}(\mu_N(p_i p_j)) \geq 2 \sqrt{\sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \frac{1}{2} \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2},$$

$$(ii) \underline{LE}(\nu_N(p_i p_j)), \overline{LE}(\nu_N(p_i p_j)) \geq 2 \sqrt{\sum_{1 \leq i < j \leq n} (\nu_N(p_i p_j))^2 + \frac{1}{2} \sum_{i=1}^n \left( d_{\nu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \nu_N(p_i p_j)}{n} \right)^2}.$$

**Proof.**

$$\begin{aligned} \left( \sum_{i=1}^n |\rho_i| \right)^2 &= \sum_{i=1}^n |\rho_i|^2 + 2 \sum_{1 \leq i < j \leq n} |\rho_i| |\rho_j| \geq 4P_\mu \\ \underline{LE}(\mu_N(p_i p_j)) &\geq 2\sqrt{P_\mu} \end{aligned}$$

Since

$$P_\mu = \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \frac{1}{2} \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2.$$

Therefore,

$$\underline{LE}(\mu_N(p_i p_j)) \geq 2 \sqrt{\sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \frac{1}{2} \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2}.$$

Similarly, we have to show that  $\overline{LE}(\mu_N(p_i p_j)), \underline{LE}(\nu_N(p_i p_j)), \overline{LE}(\nu_N(p_i p_j))$ .  $\square$

**Example 3.** Consider IFRG,  $\xi = (\underline{\xi}, \overline{\xi}) = (M, N)$  and  $V = \{p_1, p_2, p_3, p_4, p_5\}$  as shown in Figure (1). Then:

From theorem (4), we have lower approximations as follows:

$$\underline{LE}(\underline{\mu}_N(p_i p_j)) \geq 2\sqrt{1.1800 + \frac{1}{2}(0.0520)} \text{ implies } 2.7200 \geq 2.1964.$$

$$\underline{LE}(\underline{\nu}_N(p_i p_j)) \geq 2\sqrt{0.3700 + \frac{1}{2}(0.0120)} \text{ implies } 1.5200 \geq 1.2264.$$

From theorem (4), we have upper approximations as follows:

$$\overline{LE}(\overline{\mu}_N(p_i p_j)) \geq 2\sqrt{2.7600 + \frac{1}{2}(0.1080)} \text{ implies } 4.1600 \geq 3.3550.$$

$$\overline{LE}(\overline{\nu}_N(p_i p_j)) \geq 2\sqrt{0.4200 + \frac{1}{2}(0.5000)} \text{ implies } 1.9900 \geq 1.6371.$$

**Theorem 5.** Let  $\xi = (\underline{\xi}, \overline{\xi}) = (M, N)$  be an IFRG on  $n$  vertices, and let  $L(\xi) = (L(\mu_N(p_i p_j)), L(\nu_N(p_i p_j)))$  be the Laplacian matrix of  $\xi$ . Then:

$$(i) \underline{LE}(\mu_N(p_i p_j)), \overline{LE}(\mu_N(p_i p_j)) \leq |\rho_1| + \sqrt{(n-1) \left( 2 \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2 - \rho_1^2 \right)},$$

$$(ii) \underline{LE}(\nu_N(p_i p_j)), \overline{LE}(\nu_N(p_i p_j)) \leq |\sigma_1| + \sqrt{(n-1) \left( 2 \sum_{1 \leq i < j \leq n} (\nu_N(p_i p_j))^2 + \sum_{i=1}^n \left( d_{\nu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \nu_N(p_i p_j)}{n} \right)^2 - \sigma_1^2 \right)}.$$

**Proof.** Apply Cauchy-Schwarz inequality to  $(1, 1, \dots, 1)$  and  $(|\rho_1|, |\rho_2|, \dots, |\rho_n|)$ , we get:

$$\sum_{i=1}^n |\rho_i| \leq \sqrt{n \sum_{i=1}^n |\rho_i|^2}$$

$$\sum_{i=2}^n |\rho_i| \leq \sqrt{(n-1) \sum_{i=2}^n |\rho_i|^2}$$

$$L\underline{E}(\mu_N(p_i p_j)) \leq |\rho_1| + \sqrt{(n-1)(2P_\mu - \rho_1^2)}$$

Since,

$$P_\mu = \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \frac{1}{2} \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2.$$

Therefore,

$$L\underline{E}(\mu_N(p_i p_j)) \leq |\rho_1| + \sqrt{(n-1) \left( 2 \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 + \sum_{i=1}^n \left( d_{\mu_N(p_i p_j)}(p_i) - \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n} \right)^2 - \rho_1^2 \right)}.$$

Similarly, we have to prove  $L\overline{E}(\mu_N(p_i p_j))$ ,  $L\underline{E}(\nu_N(p_i p_j))$ ,  $L\overline{E}(\nu_N(p_i p_j))$ .  $\square$

**Example 4.** Consider IFRG,  $\xi = (\underline{\xi}, \overline{\xi}) = (M, N)$  and  $V = \{p_1, p_2, p_3, p_4, p_5\}$  as shown in Figure (1). Then:

From theorem (5), we have a lower approximation bound as follows:

$$L\underline{E}(\underline{\mu}_N(p_i p_j)) \leq 1.36 + \sqrt{(5-1)(2.3600 + 0.0520 - 1.8496)} \text{ implies } 2.7200 \leq 2.8599.$$

$$L\underline{E}(\underline{\nu}_N(p_i p_j)) \leq 0.76 + \sqrt{(5-1)(0.7400 + 0.0120 - 0.5776)} \text{ implies } 1.5200 \leq 1.5952.$$

From theorem (5), we have an upper approximation bound as follows:

$$L\overline{E}(\overline{\mu}_N(p_i p_j)) \leq 2.08 + \sqrt{(5-1)(5.5200 + 0.1080 - 4.3264)} \text{ implies } 4.1600 \leq 4.3618.$$

$$L\overline{E}(\overline{\nu}_N(p_i p_j)) \leq 0.40 + \sqrt{(5-1)(0.8400 + 0.5000 - 0.1600)} \text{ implies } 1.9900 \leq 2.5726.$$

**Theorem 6.** If the IFRG  $\xi = (\underline{\xi}, \overline{\xi}) = (M, N)$  is regular, then:

$$(i) L\underline{E}(\mu_N(p_i p_j)), L\overline{E}(\mu_N(p_i p_j)) \leq |\rho_1| + \sqrt{(n-1) \left( 2 \sum_{1 \leq i < j \leq n} (\mu_N(p_i p_j))^2 - \rho_1^2 \right)},$$

$$(ii) L\underline{E}(\nu_N(p_i p_j)), L\overline{E}(\nu_N(p_i p_j)) \leq |\sigma_1| + \sqrt{(n-1) \left( 2 \sum_{1 \leq i < j \leq n} (\nu_N(p_i p_j))^2 - \sigma_1^2 \right)}.$$

**Proof.** Let  $\xi$  be a regular intuitionistic fuzzy rough graph and  $d_{\mu_N(p_i p_j)}(p_i) = \frac{2 \sum_{1 \leq i < j \leq n} \mu_N(p_i p_j)}{n}$ .

Substitute this value in the theorem (5), and we get

$$LE(\underline{\mu}_N(p_i p_j)) \leq |\rho_1| + \sqrt{(n-1) \left( 2 \sum_{1 \leq i < j \leq n} (\underline{\mu}_N(p_i p_j))^2 - \rho_i^2 \right)}.$$

Similarly, we have to show that  $L\bar{E}(\underline{\mu}_N(p_i p_j)), L\underline{E}(\underline{\nu}_N(p_i p_j)), L\bar{E}(\underline{\nu}_N(p_i p_j))$ . □

**Example 5.** Consider IFRG,  $\xi = (\underline{\xi}, \bar{\xi}) = (M, N)$  and  $V = \{p_1, p_2, p_3, p_4, p_5\}$  as shown in Figure (1). Then:

From theorem (6), we have lower approximations as follows:

$$L\underline{E}(\underline{\mu}_N(p_i p_j)) \leq 1.36 + \sqrt{(5-1)(2.3600 - 1.8496)} \text{ implies } 2.7200 \leq 2.7888.$$

$$L\underline{E}(\underline{\nu}_N(p_i p_j)) \leq 0.76 + \sqrt{(5-1)(0.7400 - 0.5776)} \text{ implies } 1.5200 \leq 1.5660.$$

From theorem (6), we have upper approximations as follows:

$$L\bar{E}(\bar{\mu}_N(p_i p_j)) \leq 2.08 + \sqrt{(5-1)(5.5200 - 4.3264)} \text{ implies } 4.1600 \leq 4.2650.$$

$$L\bar{E}(\bar{\nu}_N(p_i p_j)) \leq 0.40 + \sqrt{(5-1)(0.8400 - 0.1600)} \text{ implies } 1.9900 \leq 2.0492.$$

Their Figure (2) represent the lower and upper approximation of Laplacian energy of membership and non-membership of lower bound and upper bound of an intuitionistic fuzzy rough graph.

For lower approximation membership of Laplacian energy = 2.7200. its lower bound = 2.1964 and upper bound = 3.4728.

For lower approximation non-membership of Laplacian energy = 1.5200. its lower bound = 1.2264 and upper bound = 1.9391.

For upper approximation membership of Laplacian energy = 4.1600. its lower bound = 3.3550 and upper bound = 5.3047.

For upper approximation non-membership of Laplacian energy = 1.9900. its lower bound = 1.6371 and upper bound = 2.5884.

## 5. Laplacian energy of intuitionistic fuzzy rough digraph

**Definition 10.** Let  $\xi = (\underline{\xi}, \bar{\xi}) = (M, \vec{N})$  be an intuitionistic fuzzy rough digraph (IFRDG) on  $n$  vertices. The out-degree matrix,  $D^{out}(\xi) = (D(\underline{\mu}_{\vec{N}}(p_i p_j)), D(\underline{\nu}_{\vec{N}}(p_i p_j))) = [d_{ij}]$  of  $\xi$  is a  $n \times n$  diagonal matrix defined as:  $d_{ij} = \begin{cases} d_{\xi}^{out}(p_i) & \text{if } i = j; \\ 0 & \text{otherwise.} \end{cases}$

**Definition 11.** The Laplacian matrix of an IFRDG  $\xi = (\underline{\xi}, \bar{\xi}) = (M, \vec{N})$  is defined as  $L(\xi) = (L(\underline{\mu}_{\vec{N}}(p_i p_j)), L(\underline{\nu}_{\vec{N}}(p_i p_j))) = D^{out}(\xi) - A(\xi)$ , where  $A(\xi)$  is an adjacency matrix and  $D^{out}(\xi)$  is an out degree matrix of an IFRDG of  $D$ .




Figure 2. Laplacian energy of intuitionistic fuzzy rough graph

**Definition 12.** The Laplacian matrix of spectrum of an IFRG  $L(\xi)$  is denoted by  $(R_L, S_L)$ , here  $R_L = L(\mu_{\vec{N}}(p_i p_j))$  and  $S_L = L(\nu_{\vec{N}}(p_i p_j))$  are the sets of Laplacian eigenvalues.

**Definition 13.** The Laplacian energy of an IFRG  $\xi = (\underline{\xi}, \bar{\xi}) = (M, \vec{N})$  is defined as:

$$LE(\underline{\xi}) = (LE(\mu_{\vec{N}}(p_i p_j)), LE(\nu_{\vec{N}}(p_i p_j))) = \left( \sum_{i=1}^n |\rho_i|, \sum_{i=1}^n |\sigma_i| \right),$$

$$LE(\bar{\xi}) = (LE(\bar{\mu}_{\vec{N}}(p_i p_j)), LE(\bar{\nu}_{\vec{N}}(p_i p_j))) = \left( \sum_{i=1}^n |\bar{\rho}_i|, \sum_{i=1}^n |\bar{\sigma}_i| \right).$$

Here:

$$\rho_i = \phi_i - \frac{2 \sum_{1 \leq i < j \leq n} \mu_{\vec{N}}(p_i p_j)}{n}, \quad \sigma_i = \psi_i - \frac{2 \sum_{1 \leq i < j \leq n} \nu_{\vec{N}}(p_i p_j)}{n},$$

$$\bar{\rho}_i = \bar{\phi}_i - \frac{2 \sum_{1 \leq i < j \leq n} \bar{\mu}_{\vec{N}}(p_i p_j)}{n}, \quad \bar{\sigma}_i = \bar{\psi}_i - \frac{2 \sum_{1 \leq i < j \leq n} \bar{\nu}_{\vec{N}}(p_i p_j)}{n}.$$

**Example 1.** Consider an IFRDG  $\xi = (\underline{\xi}, \bar{\xi}) = (M, \vec{N})$  on  $V = \{p_1, p_2, p_3, p_4, p_5\}$  and  $\vec{E} = \{p_1 p_2, p_2 p_5, p_5 p_4, p_4 p_3, p_3 p_1\}$ , as show in Figure (3).

From Figure (3a), the adjacency matrix, out degree matrix and Laplacian matrix of the lower IFRDG are as follows:

$$A(\underline{\xi}) = \begin{bmatrix} (0.0, 0.0) & (0.1, 0.2) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.1, 0.1) \\ (0.1, 0.2) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.1, 0.1) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.1, 0.1) & (0.0, 0.0) \end{bmatrix}$$




Figure 3. Intuitionistic fuzzy rough digraph

$$D^{out}(\underline{\xi}) = \begin{bmatrix} (0.1, 0.2) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.1, 0.1) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.1, 0.2) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.1, 0.1) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.1, 0.1) \end{bmatrix}$$

$$L(\underline{\xi}) = \begin{bmatrix} (0.1, 0.2) & (-0.1, -0.2) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.1, 0.1) & (0.0, 0.0) & (0.0, 0.0) & (-0.1, -0.1) \\ (-0.1, -0.2) & (0.0, 0.0) & (0.1, 0.2) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (-0.1, -0.1) & (0.1, 0.1) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (-0.1, -0.1) & (0.1, 0.1) \end{bmatrix}$$

The Laplacian spectrum and the Laplacian energy of an IFRDG for lower are shown in Figure (3a), as follows:

$$Laplacian\ Spec(\underline{\mu}_{\bar{N}}(p_i p_j)) = (0.0000, 0.0691+0.0951i, 0.0691-0.0951i, 0.1809+0.0588i, 0.1809-0.0588i),$$

$$Laplacian\ Spec(\underline{\nu}_{\bar{N}}(p_i p_j)) = (0.2545+0.0716i, 0.2545-0.0716i, 0.0955+0.1174i, 0.0955-0.1174i, 0.0000).$$

The calculation of the components for Laplacian energy of an intuitionistic fuzzy rough digraph for lower  $\underline{\xi}$  is  $LE(\underline{\xi}) = (0.5000, 0.6612)$ .

From Figure (3b), the adjacency matrix, out degree and Laplacian matrix of the upper IFRDG are as follows:

$$\begin{aligned}
A(\bar{\xi}) &= \begin{bmatrix} (0.0, 0.0) & (0.3, 0.1) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.2, 0.1) \\ (0.3, 0.1) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.3, 0.1) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.2, 0.1) & (0.0, 0.0) \end{bmatrix} \\
D^{out}(\bar{\xi}) &= \begin{bmatrix} (0.3, 0.1) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.2, 0.1) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.3, 0.1) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.3, 0.1) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (0.2, 0.1) \end{bmatrix} \\
L(\bar{\xi}) &= \begin{bmatrix} (0.3, 0.1) & (-0.3, -0.1) & (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.2, 0.1) & (0.0, 0.0) & (0.0, 0.0) & (-0.2, -0.1) \\ (-0.3, -0.1) & (0.0, 0.0) & (0.3, 0.1) & (0.0, 0.0) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (-0.3, -0.1) & (0.3, 0.1) & (0.0, 0.0) \\ (0.0, 0.0) & (0.0, 0.0) & (0.0, 0.0) & (-0.2, -0.1) & (0.2, 0.1) \end{bmatrix}
\end{aligned}$$

The Laplacian spectrum and the Laplacian energy of an IFRDG for upper are shown in Figure (3b), as follows:

$$Laplacian\ Spec(\bar{\mu}_{\bar{N}}(p_i p_j)) = (0.4701+0.1474i, 0.4701-0.1474i, 0.1799+0.2380i, 0.1799-0.2380i, 0.0000),$$

$$Laplacian\ Spec(\bar{\nu}_{\bar{N}}(p_i p_j)) = (0.0000, 0.0691+0.0951i, 0.0691-0.0951i, 0.1809+0.0588i, 0.1809-0.0588i).$$

The calculation of the components for Laplacian energy of an intuitionistic fuzzy rough digraph for upper  $\bar{\xi}$  is  $L\bar{E}(\bar{\xi}) = (1.2754, 0.5000)$ .

Hence, the Laplacian energy of an intuitionistic fuzzy rough digraph  $\xi$  is

$$LE(\xi) = (L\underline{E}(\xi), L\bar{E}(\bar{\xi})) = ((0.5000, 0.6612), (1.2754, 0.5000)).$$

### 5.1. Intuitionistic fuzzy rough preference relation (IFRPR)

In this part, we present a group decision-making method with intuitionistic fuzzy rough preference relation. Let  $C = \{c_1, c_2, \dots, c_n\}$  be a set of  $n$  alternatives, which is evaluated by a set  $e = \{e_1, e_2, \dots, e_m\}$  of decision-makers. Without loss of the generality, assume that the measures  $e_1, e_2, \dots, e_k$  are an advantage, while rules  $e_{k+1}, e_{k+2}, \dots, e_m$  are a disadvantage. Furthermore, let  $R_k = \left( r_{ij}^{(k)} \right)_{n \times n}$  be an intuitionistic fuzzy rough preference relation, where  $k = 1, 2, \dots, p$ . Weight  $w_r$  are assigned to the measurements  $e_r$  so that  $\sum_{r=1}^n w_r = 1$ . It is necessary to select the most optimal alternative.

For IFRGs, the cumulative grid may be obtained by intuitionistic fuzzy rough weighted averaging (IFRWA) [8], as shown below.



$$IFRWA \left( R_{ij}^{(1)}, R_{ij}^{(2)}, \dots, R_{ij}^{(n)} \right) = \left( 1 - \prod_{r=1}^s \left( 1 - \mu_{jk}^{(r)} \right)^{w_r}, \prod_{r=1}^s \left( \nu_{jk}^{(r)} \right)^{w_r} \right).$$

Where  $w_r$  be the weight function,  $R_{ij}^{(1)}, R_{ij}^{(2)}, \dots, R_{ij}^{(n)}$  be an individual intuitionistic fuzzy rough preference relation (IFRPR),  $\mu_{jk}$  be the membership value,  $\nu_{jk}$  be the non-membership value.

Here  $w_r$  are weight of  $n$  number of Laplacian energies calculating by using following formula.

For lower,

$$\underline{w}_r = \left( \left( \underline{w}_\mu \right)_r, \left( \underline{w}_\nu \right)_r \right) = \left( \frac{LE \left( \left( \underline{D}_\mu \right)_r \right)}{\sum_{i=1}^s LE \left( \underline{D}_\mu \right)_i}, \frac{LE \left( \left( \underline{D}_\nu \right)_r \right)}{\sum_{i=1}^s LE \left( \underline{D}_\nu \right)_i} \right),$$

For upper,

$$\overline{w}_r = \left( \left( \overline{w}_\mu \right)_r, \left( \overline{w}_\nu \right)_r \right) = \left( \frac{LE \left( \left( \overline{D}_\mu \right)_r \right)}{\sum_{i=1}^s LE \left( \overline{D}_\mu \right)_i}, \frac{LE \left( \left( \overline{D}_\nu \right)_r \right)}{\sum_{i=1}^s LE \left( \overline{D}_\nu \right)_i} \right),$$

$r = 1, 2, \dots, s.$

## 5.2. Algorithm

The algorithm for the selection of the most significant data visualization.

**INPUT:** A set of alternatives  $C = \{c_1, c_2, \dots, c_n\}$ , a set of experts  $e = \{e_1, e_2, \dots, e_m\}$  and build of intuitionistic fuzzy rough preference relation for lower and upper of  $R_k = \left( r_{ij}^{(k)} \right)_{n \times n}$  for each expert  $k$ .

**OUTPUT:** The process of selecting the best visualization.

1. Start

2. Determine the Laplacian energy of every IFRDG  $C_k, k = 1, 2, \dots, n.$

$$\begin{aligned} \text{For lower, } LE(\underline{\xi}) &= \left| \phi_i - \frac{2 \sum_{1 \leq i < j \leq n} \underline{\mu}_{\overline{N}}(p_i p_j)}{n} \right|, LE(\underline{\xi}) = \left| \psi_i - \frac{2 \sum_{1 \leq i < j \leq n} \underline{\nu}_{\overline{N}}(p_i p_j)}{n} \right|, \\ \text{For upper, } LE(\overline{\xi}) &= \left| \overline{\phi}_i - \frac{2 \sum_{1 \leq i < j \leq n} \overline{\mu}_{\overline{N}}(p_i p_j)}{n} \right|, LE(\overline{\xi}) = \left| \overline{\psi}_i - \frac{2 \sum_{1 \leq i < j \leq n} \overline{\nu}_{\overline{N}}(p_i p_j)}{n} \right|. \end{aligned} \quad (1)$$

3. Compute the weight vector for specialists based on the Laplacian energy of IFRDG, by utilizing

$$\begin{aligned}
\text{For lower, } \underline{w}_r &= \left( (\underline{w}_\mu)_r, (\underline{w}_\nu)_r \right) = \left( \frac{LE \left( (\underline{D}_\mu)_r \right)}{\sum_{i=1}^s LE \left( \underline{D}_\mu \right)_i}, \frac{LE \left( (\underline{D}_\nu)_r \right)}{\sum_{i=1}^s LE \left( \underline{D}_\nu \right)_i} \right), \\
\text{For upper, } \overline{w}_r &= \left( (\overline{w}_\mu)_r, (\overline{w}_\nu)_r \right) = \left( \frac{LE \left( (\overline{D}_\mu)_r \right)}{\sum_{i=1}^s LE \left( \overline{D}_\mu \right)_i}, \frac{LE \left( (\overline{D}_\nu)_r \right)}{\sum_{i=1}^s LE \left( \overline{D}_\nu \right)_i} \right).
\end{aligned} \tag{2}$$

$r = 1, 2, \dots, s.$

4. Compute the intuitionistic fuzzy rough weighted averaging by utilizing

$$IFRWA \left( R_{ij}^{(1)}, R_{ij}^{(2)}, \dots, R_{ij}^{(n)} \right) = \left( 1 - \prod_{r=1}^s \left( 1 - \mu_{jk}^{(r)} \right)^{w_r}, \prod_{r=1}^s \left( \nu_{jk}^{(r)} \right)^{w_r} \right). \tag{3}$$

5. Utilize the IFRWA operator to fuse all the individuals IFRPRs  $R_k = \left( r_{ij}^{(k)} \right)_{n \times n}$  ( $k = 1, 2, \dots, p$ ) into the collective IFRPR  $R = (r_{ij})_{n \times n}$ .

6. Calculate their scores using the score function

$$s_{ij} = \mu_{ij} - \nu_{ij}. \tag{4}$$

7. Determine the net degree of preference of visualization  $c_i$  over the other schemes by utilizing:

$$\sigma(c_i) = \sum_{r=1}^n w_r \left( \sum_{j=1, j \neq i}^n \left( r_{ij}^{(r)} - r_{ji}^{(r)} \right) \right), i = 1, 2, \dots, n. \tag{5}$$

8. Rank all the visualization  $c_i$  ( $i = 1, 2, \dots, n$ ) according to  $\sigma(c_i)$  ( $i = 1, 2, \dots, n$ ).

9. output the best data visualization.

10. Stop

## 6. Application of Laplacian energy from an IFRDG in decision-making

Group decision-making is a crucial method for identifying the best option from finite alternatives, playing a significant role in societal development and problem-solving.

This section contains, to demonstrate our proposed ideas of the intuitionistic fuzzy rough graph in decision-making, we utilise a practical example: the design of a data visualization technique.

## 6.1. The selection of data visualization for sales performance analysis

**Study case:** This section examines a case study involving the proposed Laplacian Energy of the IFRDG, applied to a real-world problem of selecting data visualization techniques for sales performance analysis. As mentioned in the introduction, data visualization for sales performance often presents a complex challenge, necessitating the selection of the most reliable strategy from various options.


The preference relation technique is extensively used to rank alternatives, with decision-makers providing their preferences regarding the available alternatives or criteria. When the information in the preference relation is expressed as intuitionistic fuzzy rough number (IFRNs), the concept of the intuitionistic fuzzy rough preference relation (IFRPR) can be redefined analogously.

**Data visualization:** Data visualization [10] may significantly improve communication and comprehension of data. When data is presented in numerical or textual representations, people may not instantly see patterns, trends, or outliers. However, effective visualization approaches may help people do just that. Visualizing data helps you comprehend and analyse it. By visualizing simplifies raw data in charts, graphs, and maps, trends, patterns, and insights may be shown. Data visualization simplifies complicated information, and clarifies results, making it crucial in corporate analysis and scientific investigation.

A company wants to analyze the sales performance of its products over the past year to make informed decisions on marketing strategies, inventory management, and product development. By employing these data visualization techniques, decision-makers can quickly and intuitively grasp the key insights from the sales data, leading to more informed decisions regarding marketing strategies, inventory management, and product development. There are four different data visualizations  $c_i$  ( $i = 1, 2, 3, 4$ ) to select from  $c_1, c_2, c_3, c_4$ . Therefore, monthly sales data for each product category, regional sales breakdown, seasonal trends, and promotions data, only the best of these would be chosen. To conduct the evaluation process based on the opinions of three independently appointed experts  $(e)_k$ , ( $k = 1, 2, 3$ ). Their comparative opinions, derived from their experience, were represented using intuitionistic fuzzy rough numbers. Additionally, intuitionistic fuzzy rough preference relations were organized into matrices as the initial step for selecting data visualization. With the proposed Laplacian energy of IFRDG utilizing these preference relations, we offer an algorithm to solve the stated visualization problem, as illustrated in Figure (4) below:

**For lower approximation:** The experts compare the involved factors among themselves and provide the initial information for computing in the form of lower intuitionistic fuzzy rough reference relations. These are represented as matrices  $R_k = (r_{ij}^{(k)})_{4 \times 4}$  ( $k = 1, 2, 3, \dots, p$ ) as shown in Figure (5) and described below:

$$\underline{R}_1 = \begin{bmatrix} (0.0, 0.0) & (0.1, 0.3) & (0.1, 0.3) & (0.1, 0.3) \\ (0.3, 0.3) & (0.0, 0.0) & (0.1, 0.3) & (0.1, 0.3) \\ (0.1, 0.3) & (0.2, 0.3) & (0.0, 0.0) & (0.1, 0.3) \\ (0.2, 0.3) & (0.2, 0.3) & (0.2, 0.3) & (0.0, 0.0) \end{bmatrix}$$



**Figure 4.** The procedure of ranking the alternatives for decision-making assessment

$$\underline{R}_2 = \begin{bmatrix} (0.0, 0.0) & (0.4, 0.2) & (0.3, 0.2) & (0.3, 0.2) \\ (0.4, 0.2) & (0.0, 0.0) & (0.3, 0.2) & (0.4, 0.2) \\ (0.3, 0.2) & (0.3, 0.2) & (0.0, 0.0) & (0.4, 0.2) \\ (0.4, 0.2) & (0.3, 0.2) & (0.4, 0.1) & (0.0, 0.0) \end{bmatrix}$$

$$\underline{R}_3 = \begin{bmatrix} (0.0, 0.0) & (0.4, 0.0) & (0.4, 0.3) & (0.5, 0.2) \\ (0.6, 0.1) & (0.0, 0.0) & (0.4, 0.3) & (0.4, 0.3) \\ (0.2, 0.1) & (0.2, 0.1) & (0.0, 0.0) & (0.1, 0.6) \\ (0.4, 0.3) & (0.5, 0.2) & (0.3, 0.4) & (0.0, 0.0) \end{bmatrix}$$

The IFRDG for lower  $C_k$  corresponding to IFRPRs in matrices  $R_k$ ,  $k = 1, 2, 3$ .

The Laplacian energy of each IFRDG for lower is calculated by using equation (1) as:

$$LE(\underline{R}_1) = (0.9000, 1.8000), LE(\underline{R}_2) = (2.6725, 1.1500), LE(\underline{R}_3) = (2.6985, 1.6191).$$




Figure 5. Intuitionistic fuzzy rough graph for lower

Then, the weight of each expert can be calculated by using equation (2) as:

$$\underline{w}_1 = (0.1435, 0.3940), \underline{w}_2 = (0.4262, 0.2517), \underline{w}_3 = (0.4303, 0.3544).$$

Hence, the experts weight vector  $e_k$   $k = 1, 2, 3$  is calculated as:

$$\underline{w} = ((0.1435, 0.3940), (0.4262, 0.2517), (0.4303, 0.3544)).$$

Utilize the aggregation operator to fuse all the individual IFRPRs  $R_k = (r_{ij}^{(k)})_{4 \times 4}$  ( $k = 1, 2, 3, 4$ ) into the collective IFRPR  $R = (r_{ij})_{4 \times 4}$ . Here, we apply the lower approximation of intuitionistic fuzzy rough weighted averaging (IFRWA) operator [8] to fuse the individual lower IFRPR. Thus, we have by using equation (3) as:

$$R = \begin{bmatrix} (0.0000, 0.0000) & (0.3641, 0.0000) & (0.3209, 0.2709) & (0.3721, 0.2346) \\ (0.4848, 0.1835) & (0.0000, 0.0000) & (0.3209, 0.2709) & (0.3641, 0.2709) \\ (0.2314, 0.1835) & (0.2443, 0.1835) & (0.0000, 0.0000) & (0.2428, 0.3463) \\ (0.3747, 0.2709) & (0.3826, 0.2346) & (0.3318, 0.2519) & (0.0000, 0.0000) \end{bmatrix}$$

Calculate their scores function by using equation (4) as:

$$s_{ij} = \begin{bmatrix} 0.0000 & 0.3641 & 0.0500 & 0.1375 \\ 0.3013 & 0.0000 & 0.0500 & 0.0932 \\ 0.0479 & 0.0608 & 0.0000 & -0.1035 \\ 0.1038 & 0.1480 & 0.0799 & 0.0000 \end{bmatrix}$$

The net flow of  $c_i$  [30], i.e., the net degree of preference of  $c_i$  over the other schemes by using equation (5). Therefore, the net flows of the four schemes are:

$$\sigma(c_1) = 0.0986, \sigma(c_2) = -0.1284, \sigma(c_3) = -0.1747, \sigma(c_4) = 0.2045.$$

Which gives the ranking of  $c_4 \succ c_1 \succ c_2 \succ c_3$ . Thus, the best visualization is  $c_4$ .

**For upper approximation:** The experts compare the involved factors among themselves and provide


the initial information for computing in the form of upper intuitionistic fuzzy rough reference relations. These are represented as matrices  $R_k = (r_{ij}^{(k)})_{4 \times 4}$  ( $k = 1, 2, 3, \dots, p$ ) as shown in Figure (6) and described below:

$$\overline{R}_1 = \begin{bmatrix} (0.0, 0.0) & (0.4, 0.1) & (0.4, 0.1) & (0.6, 0.1) \\ (0.6, 0.2) & (0.0, 0.0) & (0.5, 0.1) & (0.5, 0.1) \\ (0.6, 0.1) & (0.5, 0.2) & (0.0, 0.0) & (0.5, 0.1) \\ (0.6, 0.1) & (0.5, 0.1) & (0.6, 0.1) & (0.0, 0.0) \end{bmatrix}$$

$$\overline{R}_2 = \begin{bmatrix} (0.0, 0.0) & (0.4, 0.0) & (0.5, 0.1) & (0.6, 0.0) \\ (0.5, 0.0) & (0.0, 0.0) & (0.5, 0.3) & (0.5, 0.2) \\ (0.6, 0.0) & (0.5, 0.3) & (0.0, 0.0) & (0.6, 0.0) \\ (0.6, 0.1) & (0.5, 0.1) & (0.6, 0.1) & (0.0, 0.0) \end{bmatrix}$$

$$\overline{R}_3 = \begin{bmatrix} (0.0, 0.0) & (0.6, 0.2) & (0.3, 0.3) & (0.4, 0.1) \\ (0.7, 0.1) & (0.0, 0.0) & (0.3, 0.5) & (0.4, 0.1) \\ (0.4, 0.5) & (0.4, 0.5) & (0.0, 0.0) & (0.3, 0.5) \\ (0.4, 0.2) & (0.4, 0.2) & (0.2, 0.4) & (0.0, 0.0) \end{bmatrix}$$

The IFRDG for upper  $C_k$  corresponding to IFRPRs in matrices  $R_k$ ,  $k = 1, 2, 3$ .



**Figure 6.** Intuitionistic fuzzy rough graph for upper

The Laplacian energy of each IFRDG for upper is calculated by using equation (1) as:

$$LE(\overline{R}_1) = (3.1500, 0.7000), LE(\overline{R}_2) = (3.2000, 0.9168), LE(\overline{R}_3) = (2.4000, 2.0000).$$

Then, the weight of each expert can be calculated by using equation (2) as:

$$\overline{w}_1 = (0.3600, 0.1935), \overline{w}_2 = (0.3657, 0.2535), \overline{w}_3 = (0.2743, 0.5530).$$

Hence, the experts weight vector  $e_k$   $k = 1, 2, 3$  is calculated as:

$$\overline{w} = ((0.3600, 0.1935), (0.3657, 0.2535), (0.2743, 0.5530)).$$

Utilize the aggregation operator to fuse all the individual IFRPRs  $R_k = (r_{ij}^{(k)})_{4 \times 4}$  ( $k = 1, 2, 3, 4$ ) into the collective IFRPR  $R = (r_{ij})_{4 \times 4}$ . Here, we apply the lower approximation of intuitionistic fuzzy rough

weighted averaging (IFRWA) operator [8] to fuse the individual upper IFRPR. Thus, we have by using equation (3) as:

$$\bar{R} = \begin{bmatrix} (0.0000, 0.0000) & (0.4632, 0.0000) & (0.4145, 0.1836) & (0.5529, 0.0000) \\ (0.5989, 0.0000) & (0.0000, 0.0000) & (0.4517, 0.3217) & (0.4744, 0.1192) \\ (0.5529, 0.0000) & (0.4744, 0.3679) & (0.0000, 0.0000) & (0.4946, 0.0000) \\ (0.5529, 0.1467) & (0.4744, 0.1467) & (0.5162, 0.2152) & (0.0000, 0.0000) \end{bmatrix}$$

Calculate their scores function by using equation (4) as:

$$s_{ij} = \begin{bmatrix} 0.0000 & 0.4632 & 0.2309 & 0.5529 \\ 0.5989 & 0.0000 & 0.1300 & 0.3552 \\ 0.5529 & 0.1065 & 0.0000 & 0.4946 \\ 0.4062 & 0.3277 & 0.3010 & 0.0000 \end{bmatrix}$$

The net flow of  $c_i$  [30], i.e., the net degree of preference of  $c_i$  over the other schemes by using equation (5). Therefore, the net flows of the four schemes are:

$$\sigma(c_1) = -0.3110, \sigma(c_2) = 0.1867, \sigma(c_3) = 0.4921, \sigma(c_4) = -0.3678.$$

Which gives the ranking of  $c_3 \succ c_2 \succ c_1 \succ c_4$ . Thus, the best visualization is  $c_3$ .

## 6.2. Comparative Analysis

This section provides a comparative analysis of selected methods for evaluating data visualization, including an intuitionistic fuzzy rough geometric aggregation function chosen for comparison. The aggregation matrices used were  $R_1, R_2, R_3$  and the weights  $w_1, w_2, w_3$  as determined in equation (3) from the previous section. The aggregation was performed using the following approaches:

Intuitionistic fuzzy rough geometric aggregation operator [8]:

$$IFRGA \left( R_{ij}^{(1)}, R_{ij}^{(2)}, \dots, R_{ij}^{(n)} \right) = \left( \prod_{r=1}^s \left( \mu_{jk}^{(r)} \right)^{w_r}, 1 - \prod_{r=1}^s \left( 1 - \nu_{jk}^{(r)} \right)^{w_r} \right). \quad (6)$$

After aggregating the matrices, the resulting matrices were evaluated using the net degree approach. The evaluations for lower IFRG are  $\sigma(c_1) = -0.1175, \sigma(c_2) = -0.1497, \sigma(c_3) = -0.2440, \sigma(c_4) = 0.3194$ , ranking the alternatives as  $c_4 \succ c_1 \succ c_2 \succ c_3$ . For the upper IFRG, the evaluations were  $\sigma(c_1) = -0.1649, \sigma(c_2) = -0.1641, \sigma(c_3) = -0.0094, \sigma(c_4) = -0.1798$ , ranking order  $c_3 \succ c_2 \succ c_1 \succ c_4$ . Both aggregating orders are same order in lower and upper approximation. When the lower and upper approximation order differ, the lower approximation order is considered the best because it provides a precise approximation based on the decision criteria. As a result, alternative  $c_4$  was identified as the optimal choice, while alternative  $c_3$  received the lowest rating and was ranked last. Thus, alternatives  $c_1$  and  $c_2$  have distinct rankings.

The study results clearly indicate that the developed technique, based on Laplacian energy, is more flexible and reasonable for decision making analysis problems. In conclusion, our research introduces the

concept of the Laplacian energy of an intuitionistic fuzzy rough graph and demonstrates its application in decision-making. The findings contribute to the existing knowledge in uncertainty modelling and offer a valuable framework for addressing complex problems across various domains.

## 7. Conclusion & scope for future work

Rough set theory is a mathematical method for handling ambiguous and incomplete data. Intuitionistic fuzzy set theory addresses understanding and utilizing incomplete knowledge. Recent research suggests that integrating these two concepts can create a more expressive and adaptable framework for representing and processing incomplete information in information systems.

This paper presents a study on evaluating strategies for data visualizations. Due to the current uncertainties related to decision-makers' preferences, the intuitionistic fuzzy rough graph was chosen. The study employs the Laplacian energy of the intuitionistic fuzzy rough digraph and performs a comparative analysis using two methods for evaluating alternative strategies and four functions for aggregating intuitionistic fuzzy rough decision matrices. The results demonstrate the approach's applicability to real-world data visualization problems. The intuitionistic fuzzy rough digraph effectively captures the decision-makers' preferences, yielding credible and reliable results, unlike other directed graphs that do not account for the degree of refusal.

From the study, it is evident that the intuitionistic fuzzy rough decision matrix aggregation functions are effective. This research characterizes the Laplacian energy of an intuitionistic fuzzy rough graph using the adjacency matrix and calculates the lower and upper bounds on this energy. The Laplacian energy of the intuitionistic fuzzy rough digraph is also discussed. Future work could extend these deliberations to obtain the Laplacian energy of intuitionistic fuzzy rough hypergraphs, with appropriate comparative analyses.

## Acknowledgement

The authors are grateful to two anonymous reviewers for their valuable comments and suggestions made on the previous draft of this manuscript. This work has been supported by Vellore Institute of Technology, Vellore, Tamil Nadu, India. (VIT,Vellore).

## Funding

Shaik Noorjahan and Shaik Sharief Basha are supported by grant this research work funded was provided by Vellore Institute of Technology.

## References

- [1] AKRAM, M., ARSHAD, M., AND SHUMAIZA. Fuzzy rough graph theory with applications. *International Journal of Computational Intelligence Systems* 12, 1 (2018), 90–107.
- [2] AKRAM, M., AND ZAFAR, F. *Hybrid soft computing models applied to graph theory*, vol. 380. Springer, 2020.
- [3] ATANASSOV, K. T. *On intuitionistic fuzzy sets theory*, vol. 283. Springer, 2012.
- [4] ATANASSOV, K. T., AND ATANASSOV, K. T. *Intuitionistic fuzzy sets*. Springer, 1999.



- [5] BASHA, S. S., AND KARTHEEK, E. Laplacian energy of an intuitionistic fuzzy graph. *Indian Journal of Science and Technology* 8, 33 (2015), 1–7.
- [6] BORZECKA, H. Multi-criteria decision making using fuzzy preference relations. *Operations Research and Decisions* 22, 3 (2012), 5–21.
- [7] CHAKRABARTY, K., BISWAS, R., AND NANDA, S. Fuzziness in rough sets. *Fuzzy sets and systems* 110, 2 (2000), 247–251.
- [8] CHINRAM, R., HUSSAIN, A., MAHMOOD, T., AND ALI, M. I. Edas method for multi-criteria group decision making based on intuitionistic fuzzy rough aggregation operators. *Ieee Access* 9 (2021), 10199–10216.
- [9] CORNELIS, C., DE COCK, M., AND KERRE, E. E. Intuitionistic fuzzy rough sets: at the crossroads of imperfect knowledge. *Expert systems* 20, 5 (2003), 260–270.
- [10] DAVE SOURABH, AVANI CHHAJED, D. M. G. S. S. The art of data visualization: tools & techniques for communicating insights. *Industrial Engineering Journal* 52, 2 (2023).
- [11] DUBOIS, D., AND PRADE, H. Rough fuzzy sets and fuzzy rough sets. *International Journal of General System* 17, 2-3 (1990), 191–209.
- [12] GUTMAN, I., AND ZHOU, B. Laplacian energy of a graph. *Linear Algebra and its applications* 414, 1 (2006), 29–37.
- [13] HAQ, I. U., SHAHEEN, T., TOOR, H., SENAPATI, T., AND MOSLEM, S. Incomplete dominance-based intuitionistic fuzzy rough sets and their application in estimation of inflation rates in the least developed countries. *IEEE Access* (2023).
- [14] HE, T., CHEN, Y., AND SHI, K. Weighted rough graph and its application. In *Sixth International Conference on Intelligent Systems Design and Applications* (2006), vol. 1, IEEE, pp. 486–491.
- [15] HUSSAIN, A., MAHMOOD, T., SMARANDACHE, F., AND ASHRAF, S. Topsis approach for mcgdm based on intuitionistic fuzzy rough dombi aggregation operations. *Computational and Applied Mathematics* 42, 4 (2023), 176.
- [16] JAMIL, M. K., ANWER, S., AZEEM, M., AND GUTMAN, I. Intuitionistic fuzzy sombor indices: A novel approach for improving the performance of vaccination centers. *Communications in Combinatorics and Optimization* (2024).
- [17] KAUFMANN, A. *Introduction à la théorie des sous-ensembles flous à l'usage des ingénieurs (fuzzy sets theory)*, vol. 3. Masson, 1973.
- [18] MAHMOOD, T., AHMMAD, J., ALI, Z., AND YANG, M.-S. Confidence level aggregation operators based on intuitionistic fuzzy rough sets with application in medical diagnosis. *IEEE Access* 11 (2023), 8674–8688.
- [19] MALIK, H. M., AND AKRAM, M. A new approach based on intuitionistic fuzzy rough graphs for decision-making. *Journal of Intelligent & Fuzzy Systems* 34, 4 (2018), 2325–2342.
- [20] MAZARBHUIYA, F. A., AND SHENIFY, M. An intuitionistic fuzzy-rough set-based classification for anomaly detection. *Applied Sciences* 13, 9 (2023), 5578.
- [21] PAWLAK, Z. Rough sets. *International journal of computer and information sciences* 11 (1982), 341–356.
- [22] PENEVA, V., AND POPCHEV, I. Aggregation of fuzzy preference relations to multicriteria decision making. *Fuzzy Optimization and Decision Making* 6, 4 (2007), 351–365.
- [23] POONIA, M., AND BAJAJ, R. K. On laplacian energy of picture fuzzy graphs in site selection problem. *Journal of Intelligent & Fuzzy Systems* 41, 1 (2021), 481–498.
- [24] RAHIMI, S. S., AND FAYAZI, F. Laplacian energy of a fuzzy graph.
- [25] ROSENFELD, A. Fuzzy graphs. *Fuzzy sets and their applications to cognitive and decision process*, Elsevier (1975), 77–95.
- [26] SALAMAT, N., KAMRAN, M., ASHRAF, S., ABDULLA, M. E. M., ISMAIL, R., AND AL-SHAMIRI, M. M. Complex decision modeling framework with fairly operators and quaternion numbers under intuitionistic fuzzy rough context. *CMES-Computer Modeling in Engineering & Sciences* 139, 2 (2024), 1893–1932.
- [27] SARKAR, R. D. Pharma software—a complete overview. *International Journal of Science and Healthcare Research* 8 (2023), 166–177.
- [28] TIWARI, A., AND LOHANI, Q. D. Interval-valued intuitionistic fuzzy rough set system over a novel conflict distance measure with application to decision-making. *MethodsX* 10 (2023), 102012.
- [29] TIWARI, A. K., SAINI, R., NATH, A., SINGH, P., AND SHAH, M. A. Hybrid similarity relation based mutual information for feature selection in intuitionistic fuzzy rough framework and its applications. *Scientific Reports* 14, 1 (2024), 5958.
- [30] WANG, Y.-M., AND FAN, Z.-P. Fuzzy preference relations: Aggregation and weight determination. *Computers & Industrial Engineering* 53, 1 (2007), 163–172.
- [31] XU, W., LIU, Y., AND SUN, W. Intuitionistic fuzzy rough sets model based on  $(\theta, \varphi)$ -operators. In *2012 9th International Conference on Fuzzy Systems and Knowledge Discovery* (2012), IEEE, pp. 234–238.
- [32] ZADEH, L. A. Fuzzy sets. *Information and control* 8, 3 (1965), 338–353.
- [33] ZHAN, J., MASOOD MALIK, H., AND AKRAM, M. Novel decision-making algorithms based on intuitionistic fuzzy rough environment. *International Journal of Machine Learning and Cybernetics* 10 (2019), 1459–1485.
- [34] ZHANG, L., AND ZHU, P. Asymmetric models of intuitionistic fuzzy rough sets and their applications in decision-making. *International Journal of Machine Learning and Cybernetics* 14, 10 (2023), 3353–3380.
- [35] ZHOU, L., AND WU, W.-Z. On generalized intuitionistic fuzzy rough approximation operators. *Information Sciences* 178, 11 (2008), 2448–2465.